

# ARCHIVED DOCUMENT

UPDATED VERSION AVAILABLE ON:

<https://www.ctag-support.org.uk/antivirals/>

**Position Statement:**  
**Use of antiviral medicines for COVID-19 in adults**  
**Interim support for UK hospital clinicians**

*(This document is regularly updated.  
Please [download](#) the most recent version)*

In partnership with



## COVID-19 Therapeutics & Support Group partners

Royal College of Physicians Joint Specialty Committee for Infectious Disease (lead partner)  
Belfast Health & Social Care Trust  
Brighton and Sussex University Hospitals NHS Trust  
Cardiff and Vale University Health Board  
Guy's & St Thomas' NHS Foundation Trust  
Imperial College Healthcare  
Imperial College London  
Liverpool University Hospitals NHS Foundation Trust  
London North West University Healthcare NHS Trust  
NHS Greater Glasgow and Clyde  
Public Health England  
Royal Free London NHS Foundation Trust  
Liverpool University Hospitals NHS Foundation Trust  
Sheffield Teaching Hospital NHS Foundation Trust  
St George's University of London  
The Christie NHS Foundation Trust  
The Newcastle Upon Tyne Hospitals NHS Foundation Trust  
University College London  
University College London Hospitals NHS Foundation Trust  
University Hospital Southampton NHS Foundation Trust  
University of East Anglia  
University of Liverpool  
University of Oxford

## Document management

This document is subject to constant review. If you identify any information that needs to be updated please contact [admin.ncl-mon@nhs.uk](mailto:admin.ncl-mon@nhs.uk).

Initiated by:	Network of High Consequence Infectious Diseases (HCID)
Individuals who have overseen the development of this guidance (alphabetical order):	Content: Dr L Bell <sup>1,2</sup> , Dr J Dunning <sup>3</sup> , Dr R Gupta <sup>1</sup> , Dr Sir M Jacobs <sup>3</sup> , Dr L McCorry <sup>4</sup> , Prof M Noursadeghi <sup>1</sup> , Dr L Turtle <sup>5</sup> , Dr J Underwood <sup>6</sup> , Mr P Wade <sup>7</sup> Formulary/Governance: Mr A Barron <sup>8</sup> , Dr P Bodalia <sup>2,8</sup> , Ms M Kassam <sup>8</sup> , Dr R Sofat <sup>1,2,8</sup> <sup>1</sup> University College London <sup>2</sup> University College London Hospitals <sup>3</sup> Royal Free London <sup>4</sup> Belfast Health & Social Care Trust <sup>5</sup> University of Liverpool / Liverpool University Hospitals <sup>6</sup> Cardiff and Vale University Health Board <sup>7</sup> Guy's & St Thomas' <sup>8</sup> North Central London Joint Formulary Committee
Groups which were consulted:	COVID-19 Therapeutics Advice & Support Group (CTAG) – Antiviral subgroup
Version number:	5.0
Available on:	<a href="http://www.ctag-support.org.uk/antivirals">http://www.ctag-support.org.uk/antivirals</a>
Publication date:	2020
Review date:	This document is subject to constant review. If you identify any information that needs to be updated please contact <a href="mailto:admin.ncl-mon@nhs.uk">admin.ncl-mon@nhs.uk</a> .

## Contents

Abbreviations.....	4
Key messages.....	4
1. Aim.....	5
2. Supporting information.....	6
Membership and provenance.....	11
Document control.....	12
References.....	13
Appendix 1: Evidence base for investigational antiviral agents to treat COVID-19.....	16

ARCHIVED

## Abbreviations

Abbreviation	Meaning
CMA	Conditional Marketing Authorisation
CMO	Chief Medical Officer
CUP	Compassionate Use Programme
DHSC	Department of Health and Social Care
EMA	European Medicines Agency
MHRA	Medicines Healthcare Products Regulatory Agency
NIHR	National Institute for Health Research
R&D	Research and Development
SARS-Cov-2	Severe acute respiratory syndrome coronavirus 2
SoC	Standard of Care

### Key messages

- Remdesivir (NHS stock) is in limited supply and should only be offered to patients who meet **all** the 'eligibility criteria' and 'additional criteria' as outlined in the UK [interim Clinical Commissioning Policy](#) for remdesivir
- Co-administration of dexamethasone or hydrocortisone is recommended for patients with critical or severe COVID-19, or those requiring supplemental oxygen to maintain target saturation – see [CTAG treatment pathway](#)
- Hospitals managing COVID-19 cases should make every effort to enrol COVID-19 patients in national priority clinical trials
- Suspected side effects to medicines used in coronavirus treatment should be reported via the Yellow Card COVID-19 reporting site: <https://coronavirus-yellowcard.mhra.gov.uk/>

## 1. Aim

- 1.1. To provide interim supporting information on the appropriate use of newly licensed and investigational antiviral medicines for the treatment and prevention of COVID-19 in adults, in the hospital setting. This position statement will be updated in response to relevant developments and superseded when specific national guidance is published by UK health technology assessment bodies (e.g. National Institute for Health and Care Excellence, All Wales Medicines Strategy Group or Scottish Medicines Consortium).
- 1.2. Information contained within this position statement does not represent a 'recommendation'; however it is intended to provide support to healthcare professionals when considering available treatment options in patients with COVID-19.

ARCHIVED

## 2. Supporting information

### Remdesivir

- Remdesivir (NHS stock) is in limited supply and may only be offered to patients who meet **all** the following criteria<sup>1,21,2</sup> (see 1-page summary in [Section 3](#)):
  - hospitalised with COVID-19
  - adults, and adolescents ≥ 12 years of age and ≥ 40 kg
  - with pneumonia requiring supplemental oxygen
  - eGFR ≥ 30ml/min
  - ALT < 5 times the upper limit of normal at baseline

Additional criteria during times of limited remdesivir supplies<sup>3,4</sup>:

- Not receiving mechanical ventilation or ECMO at the time of decision to treat with remdesivir
    - *Exception:* Patients who present with an initial rapid deterioration can be considered for treatment with remdesivir
  - *Where a patient is not suitable for escalation*, a multi-disciplinary team should determine if patient would benefit from remdesivir
    - There is no current trial evidence differentiating the treatment effect of RDV according to baseline performance status. CTAG recognises that other factors, for example futility of treatment, may be taken into account by a multidisciplinary team in recommending appropriate management of severely unwell patients.
  - *Where a patient on remdesivir requires escalation*, a multi-disciplinary team should determine if remdesivir should be continued
- Co-administration of dexamethasone or hydrocortisone is recommended for patients with critical or severe COVID-19, or those requiring supplemental oxygen to maintain target saturation – see [CTAG treatment pathway](#).<sup>3,4</sup>
  - A remdesivir Compassionate Use Programme (CUP) is available for children <12 years or adolescents aged 12-17 years and weighing <40 kg with severe COVID-19 – see 1-page summary in [Section 4](#).
  - CTAG encourages enrolment into [ISARIC-CCP CRF](#) (Tier 0; no consent required). If there is limited research capacity, CTAG recommends prioritising enrolment into national priority interventional clinical trials ahead of enrolment into ISARIC-CCP CRF.
  - Ensure only patients with SARS-CoV-2 infection are treated with remdesivir.<sup>5</sup> In the absence of a confirmed diagnosis, a multidisciplinary team should have a high level of confidence that the clinical and radiological features suggest that COVID-19 is the most likely diagnosis.

### 2.1. Remdesivir is licensed for the treatment of COVID-19 in adults and adolescents with pneumonia requiring supplemental oxygen

- [ACTT-1](#) suggest remdesivir reduces time to recovery in hospitalised patients with confirmed SARS-CoV-2 infection in the subgroup of patients requiring supplemental oxygen.<sup>5</sup> See [remdesivir evidence summary](#).
- [GS-5447](#) suggests a possible benefit in clinical status, in hospitalised patients with confirmed SARS CoV-2 infection and with oxygen saturations >94% on room air; however CTAG do not feel that the small and inconsistent effect size is sufficient evidence of a clinically meaningful benefit in this ‘moderate COVID-19’ patient group.<sup>6</sup> See [remdesivir evidence summary](#).

- CTAG are of the opinion that remdesivir is unlikely to improve clinical outcome in people who appear clinically to be in the recovery phase of the illness, or those who have required mechanical ventilation or ECMO for a number of days and do not have ongoing evidence of high viral burden or ongoing viral replication, nor have advanced immunosuppression that may put them at risk of reactivation.
  - Use in pregnancy: <sup>2</sup>CTAG are of the opinion that a decision to use remdesivir should be taken by an MDT (including Obstetric and Infectious Diseases specialists) due to the uncertain fetal risk profile with treatment. More guidance will follow in a future update.
- 2.2. Other antiviral medicines are being investigated for the management of COVID-19; the current evidence-base for these medicines is summarised in [Appendix 1](#).
  - 2.3. Hospitals managing COVID-19 cases, including those treated with remdesivir, should make every effort to enrol COVID-19 patients in national priority clinical trials – refer to [Section 5](#).<sup>5-7</sup>
  - 2.4. Information for the use of dexamethasone and hydrocortisone, or investigative immunomodulators for COVID-19 (e.g. tocilizumab, sarilumab, anakinra) is available at <https://www.ctag-support.org.uk/immunomodulators>
  - 2.5. Patients may be diagnosed with COVID-19 whilst receiving investigative prophylactic treatment e.g. COPCOV trial. When such a patient would otherwise be eligible for enrolment into an interventional treatment trial, discontinuation of the prophylactic agent should be discussed with the local Principal Investigator responsible for their treatment.
  - 2.6. Suspected side effects to medicines used in coronavirus treatment should be reported via the Yellow Card COVID-19 reporting site: <https://coronavirus-yellowcard.mhra.gov.uk/>

### 3. Remdesivir summary – NHS Interim Commissioning Policy (limited supply)

Remdesivir 100 mg concentrate for solution for infusion / Remdesivir 100 mg powder for concentrate for solution for infusion (Veklury®)					
Indication <sup>1,2</sup>	Exclusion criteria <sup>1,2</sup>	Dose and duration <sup>2,8</sup>	Stopping criteria <sup>1,2,8,9</sup>	Drug specific monitoring <sup>9</sup>	Supply
<p>As per <a href="#">Interim Clinical Commissioning Policy and Supply Disruption Alert for remdesivir</a>:</p> <ul style="list-style-type: none"> <li>hospitalised with COVID-19</li> <li>adults, and adolescents ≥ 12 years of age and ≥ 40 kg</li> <li>with pneumonia requiring supplemental oxygen</li> </ul> <p>Additional criteria during supply disruption:</p> <ul style="list-style-type: none"> <li>Not receiving mechanical ventilation or ECMO at the time of decision to treat with remdesivir <ul style="list-style-type: none"> <li><i>Exception:</i> Patients who present with an initial rapid deterioration can be considered for treatment with remdesivir</li> </ul> </li> <li>Where a patient is not suitable for escalation, a MDT should determine if patient would benefit from remdesivir<sup>a</sup></li> <li>Where a patient on remdesivir requires escalation, a MDT should determine if remdesivir should be continued.</li> </ul>	<ul style="list-style-type: none"> <li>eGFR &lt; 30ml/min</li> <li>ALT ≥ 5 times the upper limit of normal (ULN) at baseline</li> </ul>	<ul style="list-style-type: none"> <li>Day 1 – single loading dose of remdesivir 200 mg given by intravenous infusion</li> <li>Day 2 to 5 – 100 mg given once daily by intravenous infusion</li> </ul> <p><b>Maximum 5 days.</b></p> <p>Infuse dose over 30 -120 minutes; see the <a href="#">SPC</a> for the relevant product for full details, and Medusa for local variations:  <a href="https://injmed.wales.nhs.uk/IVGuideDisplay.asp">https://injmed.wales.nhs.uk/IVGuideDisplay.asp</a></p>	<p>Remdesivir should be discontinued in patients who develop:</p> <ul style="list-style-type: none"> <li>ALT ≥ 5 times the ULN during treatment. It may be restarted when the ALT is &lt; 5 times the ULN</li> <li>ALT elevation accompanied by signs or symptoms of liver inflammation or increasing conjugated bilirubin, alkaline phosphatase or INR.</li> <li>eGFR &lt;30 mL/min</li> </ul> <p>Where a patient requires escalation, a MDT should determine if remdesivir should be continued.</p>	<p>Renal and liver function should be monitored carefully during treatment with remdesivir as clinically appropriate.</p>	<p>Refer to the <a href="#">Interim Clinical Commissioning Policy</a></p>

<sup>a</sup> There is no current trial evidence differentiating the treatment effect of RDV according to baseline performance status. CTAG recognises that other factors, for example futility of treatment, may be taken into account by a multidisciplinary team in recommending appropriate management of severely unwell patients.



#### 4. Remdesivir summary – Gilead compassionate use programme

Remdesivir infusion (formerly GS-5734)				
Eligibility criteria <sup>10</sup>	Exclusion criteria <sup>10</sup>	Dose and duration	Drug specific monitoring <sup>9</sup>	Supply route <sup>10</sup>
<ul style="list-style-type: none"> <li>• Children &lt;12years of age</li> <li>• Adolescents aged 12-17 years and weighing &lt;40 kg</li> </ul> <p>And</p> <ul style="list-style-type: none"> <li>• Hospitalization</li> <li>• Confirmed COVID-19</li> <li>• Severe manifestation of disease</li> </ul>	<ul style="list-style-type: none"> <li>○ Evidence of Multi-organ failure</li> <li>○ Pressor requirement to maintain blood pressure</li> <li>○ ALT levels &gt; 5 X ULN</li> <li>○ CrCl &lt;30 mL/min or dialysis or Continuous Veno-Venous Hemofiltration</li> <li>○ Concomitant administration of other investigational medicines for COVID-19 is not permitted while receiving remdesivir.</li> </ul>	<p>Please contact Gilead directly.</p>	<p>Renal and liver function should be monitored carefully during treatment with remdesivir as clinically appropriate.</p>	<p>Requests for remdesivir for individual patient use at <a href="https://rdvcu.gilead.com/">https://rdvcu.gilead.com/</a>.</p> <p>Any communication with Gilead should include <a href="mailto:UKICOVID-19@gilead.com">UKICOVID-19@gilead.com</a></p>

## 5. UK clinical studies investigating antiviral medicines

- 5.1. Organisations should prioritise support for Urgent Public Health COVID-19 studies which have been nationally prioritised. Non-prioritised research should continue, subject to it not having a negative impact on the system's ability to recruit participants and provide the resources needed to support priority clinical studies.<sup>6</sup>
- 5.2. A complete list of nationally prioritised research studies for COVID-19 is available on the NIHR website <https://www.nihr.ac.uk/covid-studies/>
- 5.3. For trials investigating immunomodulatory medicines, please refer to <https://www.ctag-support.org.uk/immunomodulators>

ARCHIVED

## Membership and provenance

The support contained within this document is provided by the COVID-19 Therapeutics Advice & Support Group (CTAG) antiviral subgroup.

The provenance for this subgroup is the Network of High Consequence Infectious Diseases (HCID). In March 2020, the collaborative expanded to include experts in Infectious Diseases from other Provider Trusts.

Organisation	Name	Role
Belfast Health & Social Care Trust	Dr Michael Hunter	Consultant, Infectious Diseases
	Dr Louise McCorry	Registrar, Infectious Diseases
Brighton and Sussex University Hospitals NHS Trust	Prof Martin Llewelyn	RCP Joint Specialty Committee for ID
Cardiff and Vale University Health Board	Dr Jonathan Underwood	Consultant, Infectious Diseases
Guys' & St Thomas' NHS Foundation Trust	Dr Anna Goodman	Consultant, Infectious Diseases
	Prof Jonathan Edgeworth	Consultant, Microbiology
	Dr Nicholas Price	Consultant, Infectious Diseases
	Dr Sam Douthwaite	Consultant, Infectious Diseases
	Dr Manu Shankar-Hari	Consultant Intensivist
	Mr Paul Wade	Consultant Pharmacist, Infectious Diseases
Imperial College Healthcare	Dr Meera Chand	Consultant, Microbiology
Imperial College London	Prof Graham Cooke	Consultant, Infectious Diseases
Imperial College London	Dr Katrina M Pollock	Clinical Research Fellow in Vaccinology
London North West University Healthcare NHS Trust	Dr Laurence John	Consultant, Infectious Diseases
Liverpool University Hospitals NHS Foundation Trust	Dr Michael Beadsworth	Consultant, Infectious Diseases
NHS Greater Glasgow and Clyde	Dr Andrew Seaton	Consultant, Infectious Diseases
Liverpool University Hospitals NHS Foundation Trust/University of Liverpool	Dr Lance Turtle	Consultant, Infectious Diseases
Royal Free London NHS Foundation Trust	Dr Jake Dunning	Consultant, Infectious Diseases
	Sir Dr Michael Jacobs	Consultant, Infectious Diseases
	Dr Sanjay Bhagani	Consultant, Infectious Diseases
Sheffield Teaching Hospitals NHS Foundation Trust	Dr Anne Tunbridge	Consultant, Infectious Diseases
	Dr Thushan de Silva	Consultant, Infectious Diseases
St George's University of London	Prof Tom Harrison	Consultant, Infectious Diseases
The Newcastle Upon Tyne Hospitals NHS Foundation Trust	Dr David Price	Consultant, Infectious Disease
	Dr Matthias Schmid	Consultant, Infectious Diseases
	Dr Yusri Taha	Consultant, Virology
University College London Hospitals NHS Foundation Trust	Dr Michael Brown	Consultant, Infectious Diseases
	Prof Mahdad Noursadeghi	Consultant, Infectious Diseases
University of Oxford	Prof Timothy Peto	Consultant, Infectious Disease

## Document control

Date	Version	Amendments
23 Mar 2020	1.0	New document
23 Mar 2020	1.1	Updated Gilead remdesivir Compassionate Use Programme eligibility criteria
27 Mar 2020	1.2	Added NHS England and NHS Improvement speciality guide for patient management. Added MHRA advice on chloroquine and hydroxychloroquine. Added hydroxychloroquine arm of RECOVERY study; added link to NIHR website.
01 Apr 2020	2.0	Updated evidence summaries in Appendix 1 (new trials for lopinavir/ritonavir and chloroquine). Merged 'Position statement' and 'Decision Support Tool' into a single document. Updated Section 3 with new trials. Updated Figure 1 with relationship between RECOVERY and PRINCIPLE/REMAP-CAP.
07 Apr 2020	2.1	Updated title to reflect updated scope (antiviral use in hospitals). Updated Section 2, 3 and Figure 1 with new trials and trial status. Included reference to CMO letter. Reformatting throughout. Added BIA and UKCPA-PIN logos (with permission).
08 Apr 2020	2.2	Updated membership & provenance. Corrected typo (2.5).
18 Apr 2020	2.3	Updated arms of RECOVERY study. Provided additional information for Gilead remdesivir Compassionate Use Programme. Updated references and web links. Updated evidence summaries in Appendix 1. Added azithromycin and inhaled interferon tables to Appendix 1.
01 May 2020	2.4	Addition of 'Key messages' and 'Abbreviations'. Emphasised importance of avoiding off-label use of investigational antivirals. Added reference to 'CTAG: Position Statement on the use of investigational immunomodulatory agents for COVID-19 in adults'. Format change to clinical trials tables. Change SNG016 and COPCOV from 'proposed' to 'active'. Added GenOMICC study. Updated evidence summaries in Appendix 1 (remdesivir; nebulised interferon; added EMA statement for chloroquine/hydroxychloroquine).
22 May 2020	2.5	Included reference to CMO letter and COVID-19 Yellow Card reporting. Updated evidence summaries in Appendix 1 (interferon). Updated support for patients diagnosed with COVID-19 whilst receiving investigative prophylactic treatment. Removed reference to GenOMICC as out of scope. Updated RECOVERY eligibility criteria to include paediatrics. Changed COPCOV from chloroquine to hydroxychloroquine. Changed CROWN CORONATION from hydroxychloroquine to chloroquine. Removed DisCoVeRY from Table 2.
06 June 2020	3.0	Included unlicensed remdesivir EAMS and NHS implementation plan for that scheme; updated Section 2 accordingly and added a new monograph (Section 5). Moved studies ACTT-1, 5773, 5774 and SNG016 into a new 'Closed to recruitment' table. Updated evidence summaries in Appendix 1 (remdesivir; chloroquine/hydroxychloroquine).
12 June 2020	3.1	Updated NHS implementation plan for EAMS. Updated evidence summaries in Appendix 1 (remdesivir; chloroquine/hydroxychloroquine).
29 June 2020	3.2	Made reference to RECOVERY dexamethasone results. Added CTAG advice on the use of remdesivir. Updated arms of PRINCIPLE study. Updated evidence summaries in Appendix 1 (systemic interferon; convalescent plasma).
03 July 2020	3.3	Moved ACCORD-2 to 'Active studies' table. Added discontinued arms of RECOVERY to 'Closed to recruitment' table. Updated evidence summaries in Appendix 1 (lopinavir/ritonavir).
07 July 2020	4	Updated with information on conditional marketing authorisation for remdesivir; EAMS programme lapsed so references removed
18 Aug 2020	4.1	Reformatting throughout. Updated evidence summaries in Appendix 1 (chloroquine/hydroxychloroquine, azithromycin, interferon (nebulised), convalescent plasma).
15 Oct 2020	5.0	Updated to reflect supply restrictions to remdesivir. Removed summary of individual trials. Removed prophylaxis chapter as out of scope. Updated all evidence summaries in Appendix 1.

## References

1. CEM/CMO/2020/035: Publication of a revised interim clinical commissioning policy: Remdesivir for patients hospitalised with COVID-19 (adults and children aged 12 years and older). Published online November 6, 2020. [https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment\\_id=103705](https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment_id=103705)
2. NHS. Interim Clinical Commissioning Policy: Remdesivir for patients hospitalised with COVID-19 (adults and children 12 years and older) Version 2. Published online November 6, 2020. [https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment\\_id=103704](https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment_id=103704)
3. Department of Health and Social Care, Medicines and Healthcare Regulatory Agency, NHS. CEM/CMO/2020/028(U) - Publication of an interim clinical commissioning policy: Remdesivir for patients hospitalised with COVID-19 (adults and children aged 12 years and older). Published online September 3, 2020. [https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment\\_id=103672](https://www.cas.mhra.gov.uk/ViewandAcknowledgment/ViewAttachment.aspx?Attachment_id=103672)
4. Powis S, Ridge K. COVID-19 therapy: corticosteroids including dexamethasone and hydrocortisone. Published online November 13, 2020. [https://www.england.nhs.uk/wp-content/uploads/2020/11/C0870\\_COVID-19-Therapy-Corticosteroids-including-dexamethasone-and-hydrocortisone-letter\\_131120.pdf](https://www.england.nhs.uk/wp-content/uploads/2020/11/C0870_COVID-19-Therapy-Corticosteroids-including-dexamethasone-and-hydrocortisone-letter_131120.pdf)
5. Atherton F, Calderwood C, McBride M, Whitty C, Powis S. CEM/CMO/2020/012 - Novel Coronavirus: Clinical Trials. [https://www.cas.mhra.gov.uk/ViewAndAcknowledgment/viewAttachment.aspx?Attachment\\_id=103495](https://www.cas.mhra.gov.uk/ViewAndAcknowledgment/viewAttachment.aspx?Attachment_id=103495)
6. Atherton F, Smith G, McBride M, Whitty C, Powis S. CEM/CMO/2020/020 - Recruiting Patients for Clinical Trials for COVID-19 Therapeutics. <https://nice.us8.list-manage.com/track/click?u=7864f766b10b8edd18f19aa56&id=230896317d&e=f70840a02d>
7. Atherton F, Smith G, McBride M, Whitty C, Powis S. CEM/CMO/2020/032 - Covid-19 Research And Treatment Trials. Published August 18, 2020. [https://www.cas.mhra.gov.uk/ViewAndAcknowledgment/viewAttachment.aspx?Attachment\\_id=103655](https://www.cas.mhra.gov.uk/ViewAndAcknowledgment/viewAttachment.aspx?Attachment_id=103655)
8. Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *The Lancet*. 2020;395(10236):1569-1578. doi:10.1016/S0140-6736(20)31022-9
9. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the Treatment of Covid-19 — Final Report. *New England Journal of Medicine*. 2020;0(0):null. doi:10.1056/NEJMoa2007764
10. Spinner CD, Gottlieb RL, Criner GJ, et al. Effect of Remdesivir vs Standard Care on Clinical Status at 11 Days in Patients With Moderate COVID-19: A Randomized Clinical Trial. *JAMA*. 2020;324(11):1048-1057. doi:10.1001/jama.2020.16349
11. Goldman JD, Lye DCB, Hui DS, et al. Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. *New England Journal of Medicine*. 2020;0(0):null. doi:10.1056/NEJMoa2015301

12. Li Y, Xie Z, Lin W. An exploratory randomized, controlled study on the efficacy and safety of lopinavir/ritonavir or arbidol treating adult patients hospitalized with mild/moderate COVID-19 (ELACOI). *Cell Press*. Published online 2020. doi:10.1016/j.medj.2020.04.001
13. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir–Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med*. Published online March 18, 2020:NEJMoa2001282. doi:10.1056/NEJMoa2001282
14. Horby PW, Mafham M, Bell JL, et al. Lopinavir–ritonavir in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *The Lancet*. 2020;0(0). doi:10.1016/S0140-6736(20)32013-4
15. Cavalcanti AB, Zampieri FG, Rosa RG, et al. Hydroxychloroquine with or without Azithromycin in Mild-to-Moderate Covid-19. *New England Journal of Medicine*. 2020;0(0):null. doi:10.1056/NEJMoa2019014
16. Furtado RHM, Berwanger O, Fonseca HA, et al. Azithromycin in addition to standard of care versus standard of care alone in the treatment of patients admitted to the hospital with severe COVID-19 in Brazil (COALITION II): a randomised clinical trial. *The Lancet*. 2020;396(10256):959-967. doi:10.1016/S0140-6736(20)31862-6
17. Li L, Zhang W, Hu Y, et al. Effect of Convalescent Plasma Therapy on Time to Clinical Improvement in Patients With Severe and Life-threatening COVID-19: A Randomized Clinical Trial. *JAMA*. Published online June 3, 2020. doi:10.1001/jama.2020.10044
18. Gharbharan A, Jordans CCE, Geurtsvankessel C. Convalescent Plasma for COVID-19. A randomized clinical trial. :16.
19. Avendano-Sola C, Ramos-Martinez A, Munez-Rubio E, et al. *Convalescent Plasma for COVID-19: A Multicenter, Randomized Clinical Trial*. Infectious Diseases (except HIV/AIDS); 2020. doi:10.1101/2020.08.26.20182444
20. Agarwal A, Mukherjee A, Kumar G, et al. *Convalescent Plasma in the Management of Moderate COVID-19 in India: An Open-Label Parallel-Arm Phase II Multicentre Randomized Controlled Trial (PLACID Trial)*. Infectious Diseases (except HIV/AIDS); 2020. doi:10.1101/2020.09.03.20187252
21. Horby P, Mafham M, Linsell L, et al. Effect of Hydroxychloroquine in Hospitalized Patients with COVID-19: Preliminary results from a multi-centre, randomized, controlled trial. *medRxiv*. Published online July 15, 2020:2020.07.15.20151852. doi:10.1101/2020.07.15.20151852
22. Skipper CP, Pastick KA, Engen NW, et al. Hydroxychloroquine in Nonhospitalized Adults With Early COVID-19. *Annals of Internal Medicine*. Published online July 16, 2020. doi:10.7326/M20-4207
23. Tang W, Cao Z, Han M, et al. Hydroxychloroquine in patients with mainly mild to moderate coronavirus disease 2019: open label, randomised controlled trial. *BMJ*. Published online May 14, 2020:m1849. doi:10.1136/bmj.m1849
24. Hung IF-N, Lung K-C, Tso EY-K, et al. Triple combination of interferon beta-1b, lopinavir–ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *The Lancet*. Published online May 2020:S0140673620310424. doi:10.1016/S0140-6736(20)31042-4

25. A Randomized Clinical Trial of the Efficacy and Safety of Interferon  $\beta$ -1a in Treatment of Severe COVID-19 | Antimicrobial Agents and Chemotherapy. Accessed October 8, 2020. <https://aac.asm.org/content/64/9/e01061-20.abstract>
26. Rahmani H, Davoudi-Monfared E, Nourian A, et al. Interferon  $\beta$ -1b in treatment of severe COVID-19: A randomized clinical trial. *International Immunopharmacology*. 2020;88:106903. doi:10.1016/j.intimp.2020.106903
27. Djukanović R, Harrison T, Johnston SL, et al. The Effect of Inhaled IFN- $\beta$  on Worsening of Asthma Symptoms Caused by Viral Infections. A Randomized Trial. *Am J Respir Crit Care Med*. 2014;190(2):145-154. doi:10.1164/rccm.201312-2235OC

ARCHIVED

# Appendix 1: Evidence base for investigational antiviral agents to treat COVID-19

## Disclaimer

Due to the urgency for interim guidance, only a limited number of agents have been assessed and a wholly systematic approach to assessing the evidence (such as GRADE) has not been performed. Some subjective judgments are solely the consensus opinion of the authors and consulted experts.

The focus here is on investigational antiviral treatments for managing hospitalised COVID-19 patients. Supportive care and treatment of co-infections and complications, such as ARDS, are not addressed.

## Methods

COVID-19 is caused by infection with the newly emerged betacoronavirus SARS-CoV-2.

We reviewed the available data on treatment of betacoronaviruses but restricted the search to investigational antiviral agents being used, or considered, within the context of UK clinical trials. This includes all investigational antiviral agents identified in the [NIHR list of nationally prioritised studies](#).

We broadly hierarchised the evidence according to the following matrix and considered the available safety data.

## Evidence summary

Summaries are provided for investigational antiviral treatments being used, or considered, within the context of UK clinical trials.

The summaries are divided into two categories in the following tables based on current evidence:

- Table 1: Benefit may exceed risk
- Table 2: Inadequate data to recommend use



**Table 1: Evidence base for specific therapies for SARS-CoV-2 infection: Benefit exceeds risk**

RCT = randomised-controlled trial; CI = confidence interval; HCQ = hydroxychloroquine. OR = odds ratio; HR = hazard ratio.

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
Remdesivir	<p>The highest level of evidence is: 4 RCTs.</p> <p><b>NICE have published rapid evidence summary for remdesivir:</b> <a href="https://www.nice.org.uk/advice/es27/chapter/Key-messages">https://www.nice.org.uk/advice/es27/chapter/Key-messages</a></p> <p><b>BMJ have published a rapid evidence summary for remdesivir:</b> <a href="https://www.bmj.com/content/370/bmj.m2924">https://www.bmj.com/content/370/bmj.m2924</a></p> <p>A double-blinded RCT in China compared remdesivir to placebo among 237 adults with severe COVID-19 (defined as radiologically-confirmed pneumonia and either SpO<sub>2</sub> ≤94 % on air or PaO<sub>2</sub>/FiO<sub>2</sub> ratio ≤300), ≤12 days from symptom onset to enrolment<sup>8</sup>. At 28 days, remdesivir was not associated with difference in time to clinical improvement (hazard ratio 1.23; 95% CI 0.87–1.75) or overall mortality (22 (14%) died in the remdesivir group vs 10 (13%) in the placebo group). In a post-hoc analysis, patients receiving remdesivir had a numerically faster time to clinical improvement than those receiving placebo among patients with symptom duration ≤10 days at enrolment (hazard ratio 1.52; 0.95–2.43). However, the trial was underpowered.</p> <p>A double-blind, randomized, controlled trial among adults hospitalized with COVID-19 with evidence of lower respiratory tract involvement (radiographic infiltrates, peripheral oxygen saturation ≤94% on room air, or requiring supplemental oxygen, mechanical ventilation, or extracorporeal membrane oxygenation) compared remdesivir to placebo.<sup>9</sup> A total of 903/1,062 (85%) participants had severe disease (SpO<sub>2</sub> ≤ 94% on room air, respiratory rate ≥24/min, requiring oxygen, invasive or non-invasive ventilation) at enrolment. Participants receiving remdesivir had a median recovery time of 10 days (95% CI 9 to 11), vs. 15 days (95% CI 13 to 18) in the placebo arm (rate ratio for recovery 1.29; 95% CI 1.12 to 1.49; log rank p&lt;0.001). The rate ratio for recovery was largest among patients with a baseline ordinal score of 5 (receiving low flow oxygen therapy; rate ratio for recovery 1.45; 95% CI 1.18 to 1.79). The benefit of remdesivir was also larger when given earlier in the illness. Patients who underwent randomization during the first 10 days of illness had a rate ratio for recovery of 1.37 (95% CI 1.14 to 1.64), compared to 1.20 (95% CI 0.94 to 1.52) &gt;10 days after symptom onset. Mortality was numerically lower in the remdesivir group than in the placebo group, but the difference was not significant. Kaplan Meier 15- and 29-day mortality estimates were 6.7% with remdesivir vs. 11.9% with placebo and 11.4% with remdesivir vs. 15.2% with placebo, respectively (hazard ratio, 0.73; 95% CI 0.52 to 1.03).</p> <p>An open-label RCT compared the efficacy of 5 vs. 10 days of remdesivir treatment vs. standard care on clinical status (7-point ordinal scale) on day 11 after treatment initiation among 596 patients admitted with moderate COVID-19 pneumonia (pulmonary infiltrates and room-air oxygen saturation &gt;94%)<sup>10</sup>. On day 11, patients in the 5-day remdesivir group had higher odds of better clinical status than those receiving standard care (odds ratio, 1.65; 95% CI, 1.09-2.48; P = .02), but clinical status was not significantly different between the 10-day remdesivir and standard care groups (P = .18). At day 28, 2 patients (1%) in the 5-day remdesivir group died, compared to 3 (2%) in the 10-day remdesivir group, and 4 (2%) in the standard care group.</p> <p>An open-label RCT of 397 hospitalized patients with SARS-CoV-2 pneumonia compared 10-day vs. 5-day courses of remdesivir.<sup>11</sup> By day 14, clinical improvement occurred in 64% of patients in the 5-day group and in 54% in the 10-day group. After adjustment for baseline clinical status, there was no significant difference between 5-day and 10-day courses at day 14 (P=0.14).</p> <p>Results from a ‘living’ meta-analysis for these summaries is available <a href="#">on the Cochrane website</a>.</p>	<p>See <a href="#">SPC</a> &amp; <a href="#">BNF</a>.</p> <p>No significant adverse safety signals detected in the COVID-19 RCTs.</p>	<p>Refer to the <a href="#">Interim Clinical Commissioning Policy</a> and the compassionate use programme criteria at: <a href="https://rdvcu.gilead.com">https://rdvcu.gilead.com</a></p>

**Table 2. Evidence base for specific therapies for SARS-CoV-2 infection: Inadequate data to recommend use**

RCT = randomised-controlled trial; CI = confidence interval; HCQ = hydroxychloroquine. OR = odds ratio; HR = hazard ratio.

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
<p><b>Lopinavir/ritonavir</b></p>	<p>The highest level of evidence is: 4 RCTs.</p> <p>An exploratory RCT assessing lopinavir/ritonavir or Arbidol® (umifenovir) to among 86 hospitalised adults with mild/moderate COVID-19 reported no differences between arms in time from positive-to-negative viral conversion using RT-PCR (mean 9.0 days (SD 5.0) in the lopinavir/ritonavir group, 9.1 (SD 4.4) in the arbidol group and 9.3 (SD 5.2) in the control group), though the trial was underpowered.<sup>12</sup></p> <p>An open-label RCT of hospitalised adults in China with severe COVID-19 (n=199) found no benefit in time to clinical improvement for lopinavir-ritonavir over standard care (hazard ratio 1.31; 95% CI 0.95 to 1.80). 28-day mortality was similar in the lopinavir-ritonavir group and the standard-care groups (19.2% vs. 25.0%; difference -5.8; 95% CI, -17.3 to 5.7). Lopinavir-ritonavir recipients spent less time in hospital (12 vs. 14 days) and less time in intensive care (6 vs. 11 days).<sup>13(p19)</sup></p> <p>The open-label RECOVERY RCT allocated 1616 patients to receive lopinavir-ritonavir, compared to 3424 patients to receive usual care.<sup>14</sup> A total of 374 (23%) patients allocated to lopinavir-ritonavir and 767 (22%) patients allocated to usual care died within 28 days (rate ratio 1.03, 95% CI 0.91-1.17; p=0.60). Results were consistent among pre-specified subgroups and for secondary endpoints of time until discharge alive from hospital (median 11 days [IQR 5 to &gt;28] in both groups), proportion of patients discharged from hospital alive within 28 days (rate ratio 0.98, 95% CI 0.91-1.05; p=0.53) and proportion who met the composite endpoint of invasive mechanical ventilation or death (risk ratio 1.09, 95% CI 0.99-1.20; p=0.092).</p> <p>The SOLIDARITY trial <a href="#">issued a press release</a> announcing closure of the lopinavir/ritonavir arm following review of interim results, and evidence from all trials presented at a WHO Summit. Lopinavir/ritonavir was noted to produce little or no reduction in the mortality of hospitalized COVID-19 patients when compared to standard care. Full results awaited.</p> <p>Results from a 'living' meta-analysis for these summaries is available <a href="#">on the Cochrane website</a>.</p>	<p>See <a href="#">SPC</a> &amp; <a href="#">BNF</a>.</p> <p>Well established agent with well understood toxicity profile. Gastrointestinal side effects are very common.</p> <p>Note multiple, significant drug-drug interactions.</p> <p>No additional significant adverse safety signals detected in the COVID-19 RCTs.</p>	<p>Licensed for the treatment of HIV-1 infection.</p> <p>Included in <a href="#">REMAP-CAP trial</a></p>

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
<b>Azithromycin</b>	<p>The highest level of evidence is: 2 RCTs.</p> <p>A multi-centre open-label RCT in Brazil (COALITION 1) compared HCQ (n=221), to HCQ plus azithromycin (n=217), to standard care (n=227) among hospitalised patients with mild-moderate COVID-19 (maximum O2 requirement of 4L/min or FiO2 40%).<sup>15</sup> In the primary modified intention to treat (mITT) analysis, there were no between-group differences in the primary outcome of clinical status at day 15 (measured on a 7 point ordinal scale; hydroxychloroquine plus azithromycin vs. standard care odds ratio 0.99; 95% CI 0.57 to 1.73; P=1.00) or in secondary outcomes including requirement for mechanical ventilation. There was low mortality in the cohort (n=18) with no between-group differences. Higher rates of adverse events were reported in the arms containing HCQ (39.3% among HCQ plus azithromycin arm; 33.7% among HCQ arm; 22.6% among standard care arm), including QTc prolongation and deranged liver function.</p> <p>A second multi-centre open-label RCT in Brazil (COALITION 2) compared azithromycin to standard of care (including HCQ therapy) among patients hospitalised patients with severe COVID-19 (oxygen &gt;4 L/min flow, high-flow nasal cannula oxygen, non-invasive mechanical ventilation or invasive mechanical ventilation).<sup>16</sup> In the primary analysis among the mITT population (n=397), there was no difference in the primary endpoint of clinical status at day 15 after randomisation (assessed by a 6-point ordinal scale) between the azithromycin and control groups (OR 1.36 [95% CI 0.94–1.97], p=0.11).</p> <p>Results from a ‘living’ meta-analysis for these summaries is available <a href="#">on the Cochrane website</a>.</p>	<p>See <a href="#">SPC</a> &amp; <a href="#">BNE</a>.</p> <p>Well established agent with well understood toxicity profile including gastrointestinal upset (common) and QT prolongation (uncommon).</p> <p>No additional significant adverse safety signals detected in the COVID-19 RCTs.</p>	<p>Various licensed indications as an antimicrobial.</p> <p>Included as an arm in the UK <a href="#">RECOVERY trial</a>.</p> <p>Prolonged macrolide therapy is also an existing arm in <a href="#">REMAP-CAP trial</a>, but with immunomodulatory rather than antiviral intent.</p>

ARCHIVED

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
<b>Convalescent plasma</b>	<p>The highest level of evidence is: 4 RCTs.</p> <p>An open-label, multicenter, randomized clinical trial in Wuhan, China compared convalescent plasma to standard care among 103 patients with severe (respiratory distress and/or hypoxemia) or life-threatening (shock, organ failure, or requiring mechanical ventilation) COVID-19<sup>17</sup>. At 28 days, there was no difference between the convalescent plasma and standard care arms in clinical improvement (51.9% vs 43.1%; hazard ratio 1.40; 95% CI 0.79-2.49; P = .26) or mortality (15.7% vs 24.0%; OR 0.65; 95% CI 0.29-1.46]; P = .30). Convalescent plasma treatment was associated with negative conversion of viral PCR at 72 hours in 87.2% of the convalescent plasma group vs 37.5% of the control group (OR 11.39; 95% CI 3.91-33.18; P &lt; .001). However, the trial was underpowered for clinical outcomes and median time between the onset of symptoms and randomization was 30 days, suggesting late initiation of therapy. Adverse events were reported for the intervention group only (see ‘Safety profile’). Furthermore, a greater proportion of participants in the treatment group received co-interventions, compared to the control group, which may have been influenced by knowledge of allocation.</p> <p>A preprint of an open-label randomized trial comparing convalescent plasma with standard of care in patients hospitalized for COVID-19 in the Netherlands reported that the trial was stopped early (n=86 enrolled) as 53 of 66 patients tested had anti-SARS-CoV-2 antibodies at baseline.<sup>18</sup> At the time of cessation, the adjusted odds ratio for overall mortality for patients treated with convalescent plasma was 0.95 (CI 0.20 – 4.67; p=0.95) and for improvement in the WHO COVID-19 disease severity score on day 15 was 1.30 (CI 0.52 - 3.32).</p> <p>A preprint of an open-label randomized trial comparing convalescent plasma with standard of care in patients hospitalized for COVID-19 in Spain was stopped after recruitment of 81 patients due to a fall in recruitment following population-level pandemic suppression.<sup>19</sup> The original intended primary endpoint was the proportion of patients in categories 5-7 of the COVID-19 ordinal scale at day 15. At enrolment, 49.4% of participants had evidence of anti-SARS-CoV-2 IgG antibodies. At closure, 0/38 patients had progressed to mechanical ventilation or death among patients assigned to receive plasma, vs. 6/43 (14%) in control arm (p=0.57). Mortality risk was 0% vs 9.3% at day 15 for the plasma and control groups, respectively. However, the study was severely underpowered due to early closure.</p> <p>A preprint of an open-label randomized trial in India compared convalescent plasma with standard of care among 464 hospitalized patients with moderate PCR-confirmed COVID-19 (PaO<sub>2</sub>/FiO<sub>2</sub> ratio: 200-300mmHg; or respiratory rate &gt; 24/min and SpO<sub>2</sub> ≤ 93% on room air).<sup>20</sup> There was no difference in the composite primary outcome of progression to severe disease (PaO<sub>2</sub>/FiO<sub>2</sub>&lt;100 mmHg; 44/235 (18.7%) in intervention arm vs. 41/229 (17.9%) in the control arm; OR 1.09; 95% CI 0.67 - 1.77]) or mortality (34/235 (13.6%) vs. 31/229 (14.6%); OR 1.06; 95% CI: -0.61 - 1.83) at 28 days post-enrolment. Neutralising antibody titres were measured in 418 trial participants; 348 (83.2%) had detectable neutralising antibodies at enrolment.</p> <p>Results from a ‘living’ meta-analysis for these summaries is available <a href="#">on the Cochrane website</a>.</p>	<p>Transfusion-related adverse events well-recognised.</p> <p>2/52 patients who received convalescent plasma in the RCT from China<sup>17</sup> experienced transfusion-associated adverse events; both improved with supportive care. No other additional significant adverse safety signals detected in the COVID-19 RCTs.</p>	<p>Included as an arm in the UK <a href="#">RECOVERY trial</a>.</p>

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
<p><b>Hydroxychloroquine (HCQ) / Chloroquine (CQ)</b></p>	<p>The highest level of evidence is: 11 RCTs. The highest quality four trial are summarised here, in addition to the SOLIDARITY press release.</p> <p>The RECOVERY trial, a multi-arm multi-centre open-label RCT of hospitalised COVID-19 patients in the UK, has published results of its HCQ arm.<sup>21</sup> The authors report no significant difference between the HCQ arm (1561 patients) and the SoC arm (3155 patients) in the primary endpoint of all-cause mortality at 28 days: 27% mortality in the HCQ group versus 25% in the SoC group, RR 1.09 (95% CI 0.97-1.23). This finding is consistent in multiple pre-specified subgroup analyses including age and days since symptom onset. For secondary outcomes, HCQ was associated with a lower probability of discharge alive at 28 days (RR 0.90, 95% CI 0.83-0.98) and higher rate of progression to a composite outcome of death or invasive mechanical ventilation: 30.7% versus 26.9% (RR 1.14, 95% CI 1.03-1.27).</p> <p>The COALITION 1 trial, a multi-centre open-label RCT in Brazil, compared HCQ, to HCQ plus azithromycin, to SoC.<sup>15</sup> Please see the azithromycin table above for a summary of this trial.</p> <p>A double-blind RCT compared HCQ (n=212) to placebo (n=211) for treatment of non-hospitalised adults with either laboratory-confirmed COVID-19 or highly suspected COVID-19 following a high risk exposure, commenced within 4 days of symptom onset.<sup>22</sup> There was no difference in the primary endpoint of change in symptom severity at 14 days (measured on a 10 point visual analogue scale): between-group difference of -0.27 points (95% CI -0.61-0.07). The study was underpowered to detect differences in hospitalisation or mortality, but notes that the incidence of these events did not differ between groups (HCQ – 5 events, placebo – 10 events, p=0.29).</p> <p>An open-label multi-centre RCT from China (n=150) compared HCQ with SoC in hospitalised patients with mild-moderate COVID-19.<sup>23</sup> The authors reported no difference between arms in SARS-CoV-2 negative conversion in respiratory tract specimens by 28 days: 85.4% (95% CI 73.8-93.8%) vs 81.3% (71.2-89.6%). There were higher rates of adverse events (30% versus 9%) in the HCQ arm. Clinical outcomes are not reported.</p> <p>The SOLIDARITY RCT <a href="#">issued a press release</a> announcing closure of the HCQ arm following review of interim results, and evidence from all trials presented at a WHO Summit. HCQ was noted to produce little or no reduction in the mortality of hospitalized COVID-19 patients when compared with SoC. Publication of the full results is awaited.</p> <p>Results from a ‘living’ meta-analysis for these summaries is available <a href="#">on the Cochrane website</a>.</p>	<p>See <a href="#">SPC</a> &amp; <a href="#">BNE</a>.</p> <p>Well established agent, defined safety profile as antimalarial drug; however, safety in acute viral illness is not established and <a href="#">concerns about the risk of QTc prolongation</a> were raised following early observational trials.</p> <p>The COALITION 1 trial reported more frequent adverse events in participants in HCQ-containing arms.<sup>15</sup> These included QTc prolongation in 30 of 438 patients who received HCQ (in comparison to 1 of 227 who did not), and elevated liver transaminases in 43 of 438 (in comparison to 8 of 227).</p> <p>The RECOVERY trial<sup>21</sup> reports that the excess risk of death seen in its HCQ group was not due to mortality due to COVID-19 (24.0% vs 23.5% in the SoC group), but notes a small excess risk of death from cardiac causes (mean±SE excess, 0.4±0.2 percentage points) and non-SARS-CoV-2 infection (mean excess, 0.4±0.2 percentage points). There were no differences in the overall frequency of arrhythmia in the participants for whom this data was available in the HCQ arm (735 patients) or the SoC arm (1421 patients). One serious adverse event attributed to HCQ was reported (torsades de pointes).</p> <p>Two RCTs have reported higher rates of adverse effects (primarily gastrointestinal) in participants receiving HCQ.<sup>22,23</sup></p>	<p>Various licensed indications, including malaria and rheumatoid arthritis.</p> <p>Included in <a href="#">COP-COV trial</a> (prophylaxis)</p>

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
<b>Interferon (systemic)</b>	<p>The highest level of evidence is: 3 RCTs.</p> <p>An open-label multicentre RCT in Hong Kong compared treatment with subcutaneous IFN<math>\beta</math>-1b, ribavirin &amp; lopinavir/ritonavir (n=86) to lopinavir/ritonavir alone (n=41), in hospitalized patients with mild-moderate COVID-19.<sup>24</sup> Patients in the group who received IFN had significantly shorter times to positive-to-negative viral conversion of NP swabs: 7 days versus 12 days, HR 4.37 (95% CI 1.86-10.24). Significant findings are also reported for secondary outcomes including time to clinical recovery in the IFN group (4 days versus 8 days) and shorter length of hospital stay (9 days versus 14.5 days). The majority of patients had mild disease and no mortality was observed. There is heterogeneity in the intervention arm, as IFN<math>\beta</math> treatment was only used in patients randomized to this group if they were within 1 week of symptom onset. A subgroup analysis indicated that the apparent benefits in the intervention group were only seen in the group of patients treated with IFN<math>\beta</math> within the first week of symptoms.</p> <p>A single-centre open-label RCT from Iran compared subcutaneous IFN<math>\beta</math>-1a (three times per week for 2 weeks, n=42) plus SoC, to SoC alone (n=39), in patients with severe COVID-19.<sup>25</sup> SoC comprised HCQ and lopinavir/ritonavir for all patients, and antibiotics and corticosteroids in some patients. There was no difference between groups in the primary outcome of time to clinical improvement. The authors report a significant difference in 28 day mortality as secondary outcome (19% in the IFN<math>\beta</math>-1a group vs 43.6% in the SoC group), however there is a high risk of bias due to missing outcome data as multiple patients were excluded from the intervention arm because of mortality within the 1<sup>st</sup> week of treatment.</p> <p>A second open-label RCT, from the same single centre in Iran, compared subcutaneous IFN<math>\beta</math>-1b given every 2<sup>nd</sup> day for 2 weeks (n=33) plus SoC, to SoC alone (n=33), in patients with severe COVID-19.<sup>26</sup> SoC again comprised HCQ and lopinavir/ritonavir. The authors report a significant difference in the primary outcome of time to clinical improvement: 9 days (IQR 6–10) versus 11 (9–15) days (p=0.002). Differences in secondary outcomes including discharge at day 14 are also reported: 78.79% in the IFN group versus 54.55% in the control group (OR 3.09; 95% CI: 1.05–9.11). No significant difference is reported for mortality at 28 days.</p> <p>Results from a ‘living’ meta-analysis for these summaries is available <a href="#">on the Cochrane website</a>.</p>	<p>See <a href="#">SPC</a> &amp; <a href="#">BNF</a>.</p> <p>Well established agent with defined but complex safety profile. Clinicians experienced in managing side effects should be consulted where there are concerns e.g. those who have treated hepatitis C virus (HCV) infection and multiple sclerosis.</p> <p>Known IFN-related adverse effects including flu-like symptoms and neuropsychiatric effects have been reported in IFN arms of trials, but no increased numbers of serious adverse events.<sup>25(p19)</sup></p>	<p>Several different interferons are available for systemic administration, for different licensed indications. There are insufficient data to strongly recommend a particular preparation, although IFN-<math>\beta</math> appears more promising based on available data.</p> <p>IFN-<math>\beta</math> injection: included as an arm in the immune modulation domain of <a href="#">REMAP-CAP trial</a> (recruiting)</p>

Therapy	Data: SARS-CoV-2	Safety profile	UK feasibility
<b>Interferon (nebulised)</b>	<p>The highest level of evidence is: 1 RCTs (data press-released but not yet published).</p> <p>The SNG016 trial is a phase II double-blind multi-centre RCT of SNG001 (a nebulized formulation of IFN<math>\beta</math>-1a) versus placebo in hospitalized patients with COVID-19, run by Synairgen. The trial issued results as <a href="#">a press release in July 2020</a> and <a href="#">further interim analyses in September 2020</a>. The more detailed interim analysis reports that patients who received the drug (n=48) had significantly greater odds of clinical improvement by day 15/16 of illness as assessed on an 8 point ordinal scale, in comparison to the placebo group (n=50): OR 2.32 (95% CI 1.07-5.04). There were no significant differences between the groups in the time to development of severe disease or death (HR 0.50, 95% CI 0.18-1.38), or in the odds of severe disease or death (OR 0.28, 95% CI 0.07-1.08), when analysed in the intention to treat population. A significant difference is reported in the per-protocol population (n=86) for the odds of severe disease or death (OR 0.15, 95% CI 0.04-0.93). The event rates in each group were not reported and overall mortality in the trial was low (3/98 patients, all in the placebo group). Formal publication of the full results is required for full interpretation. The company <a href="#">subsequently announced</a> they were extending the SNG-16 trial to a second phase recruiting patients in the community.</p>	<p>Nebulised IFN formulations not yet licensed in the UK. Please see IFN (systemic) table for medicines information for injectable IFNs.</p> <p>A phase II human trial of SNG001 (nebulized IFN <math>\beta</math>-1a), in individuals with a background of viral-induced asthma who had new cold-like symptoms, reported that it was well tolerated with no safety signals flagged.<sup>27</sup></p>	<p>Clinical formulation made by Synairgen, SNG001, a nebulised formulation of IFN<math>\beta</math>-1a. Synairgen has announced a SNG001 <a href="#">Managed Access Program</a> jointly with Clinigen, for hospitalised COVID-19 patients in Europe.</p> <p>Recruitment to the <a href="#">second community-based phase of the SNG016 trial</a>, run by Synairgen and the University of Southampton, is ongoing.</p>

ARCHIVED